

Is Your Bottled Water KILLING YOU?

By William Davis, MD

Based on the poor quality of what comes out of the municipal tap, health-conscious people often rely on bottled or home-filtered water for their drinking needs. While most filtered and bottled waters are free of cancer-causing contaminants, they provide little or no *magnesium*. Even most tap water is devoid of this critical mineral.

The implications of this widespread magnesium deficiency are frightening, in as much as communities with *low* magnesium content in drinking water show *increased* rates of sudden death.

Magnesium plays hundreds of crucial roles in the body, including suppressing unstable heart rhythms, controlling blood pressure, maintaining insulin sensitivity, and regulating over 300 enzymes. Attaining optimal magnesium levels is an absolute requirement for good health.

In this article, we examine magnesium's importance to human health, how changes in the way we obtain our drinking water have contributed to widespread magnesium deficiencies, and strategies you can use to optimize your magnesium intake through dietary sources, better bottled waters, and nutritional supplementation. > > >

Magnesium: Lost in the Filter

Our human ancestors evolved in a world in which healthy drinking water came directly from streams, rivers, and lakes, rich in mineral content. The human body became reliant on obtaining a considerable proportion of its daily mineral needs from natural water sources.

Fast-forward to the twenty-first century. We obtain drinking water from a spigot or a plastic bottle. Pesticides and other chemicals seeping into the water supply have made everyone suspicious of water quality. As a result, municipal water-purification facilities have intensified their efforts to remove contaminants like lead, pesticide residues, and nitrates from drinking water. Unfortunately, these modern water-treatment methods also deplete drinking water of *desirable* minerals like calcium and magnesium.

Exacerbating this problem is that many Americans, distrustful of the purity and safety of municipally treated water, have added home water filters and purifiers that efficiently extract any remaining minerals from the water, thus converting "hard" into "soft" water. In fact, the

manufacturers of these devices boast of their power to yield water free of "contaminants"—including minerals like magnesium. Thus, the magnesium content of the water that passes through most commercial filters is zero.¹

The present-day enthusiasm for bottled water has further compounded the problem. Americans consumed nearly *8 billion gallons* of bottled water last year. The mineral content of these products varies widely. While some mineral waters, particularly those from Europe, contain a moderate amount of magnesium, other brands of bottled water contain little or none.²

The upshot of all this is that we cannot rely on drinking water to provide adequate magnesium. The recommended dietary allowance (RDA) for magnesium—that is, the amount required to prevent *severe deficiency*—is 420 mg a day for men and 320 mg a day for women. In cities with the *highest* magnesium water content, only 30% of the RDA can be obtained by drinking *two liters* of tap water a day.⁴ In most cities, only a meager 10–20% of the daily requirement can be obtained. That leaves 70–90% of the daily magnesium requirement that

MAGNESIUM BASICS

Magnesium is required for the proper function of approximately 300 enzymes in the human body. Functions as diverse as blood pressure regulation, muscle contraction, heart rhythm stabilization, and nervous system communication are all magnesium-dependent processes. Humans cannot survive without magnesium. Some authorities have even argued that some typical manifestations of aging—such as loss of muscle mass, rising blood pressure, and diminished nervous system function—are partly attributable to magnesium deficiency.⁴

must be obtained from other sources. Since many people's diets are also low in magnesium, the average American ingests substantially less magnesium than the RDA.³


The problem may be even worse than it appears. Many authorities believe that higher levels of magnesium are needed to avoid serious illnesses such as heart disease.⁴ Others argue that "normal" magnesium blood levels reported by laboratories, originally derived from populations symptomatic with magnesium deficiency, are also too low and that higher blood levels are necessary for optimal health.⁵

Low Magnesium Tied to Risk of Sudden Death

"Results from the early epidemiological studies suggest that sudden-death rates in soft-water areas are at least 10% greater than sudden-death rates in hard-water areas.

*If magnesium supplementation causes even a modest decrease in sudden-death rates, a substantial number of lives might be saved."*⁷

—Mark J. Eisenberg, MD, MPH
McGill University



Magnesium deficiency may have potentially dire consequences. Critically ill patients, who often have very low magnesium levels, suffer the risk of seizures, life-endangering heart rhythms, and fatal heart attack.⁸

Magnesium has a stabilizing effect on cell membranes, particularly in heart muscle. A healthy heart generates stable, predictable electrical impulses. Lack of magnesium permits unstable electrical impulses in the heart to emerge, generating abnormal heart rhythms.^{9,10} In fact, much magnesium research over the years has focused on its administration during heart attack to reduce death from fatal heart rhythms.¹¹

Magnesium blood levels are routinely monitored in hospitalized patients, particularly those who are receiving diuretic medications. It is well known in hospitals that if magnesium blood levels drop too low, abnormal heart rhythms can suddenly develop, necessitating intravenous replenishment of magnesium.⁹ Dangerous rhythms of the heart's ventricles, in particular, can occur. People suffering from congestive heart failure are especially susceptible to dangerous heart rhythms when magnesium is low.

Scientists have observed that people in areas with higher levels of magnesium in their drinking water

exhibit rates of sudden cardiac death that are three to four times *lower* than those of people living in municipalities with the lowest magnesium levels in drinking water.^{7,12,13} This has drawn the attention of national and international public health officials. For example, a recent World Health Organization (WHO) report on the quality of drinking water cited 80 studies that have examined the relationship between cardiovascular death and water "hardness" (measured principally by magnesium and calcium content). The WHO concluded that the magnesium content of water is indeed a cardiovascular risk factor and that supplementing drinking water with magnesium should be a priority, much as fluoride became one.¹⁴ To date, however, no action has been taken.

The Many Perils of Magnesium Deficiency

The older you are, the more likely magnesium depletion is to develop,¹⁵ with substantial deficiency common by the age of 50. If you have any condition that causes frequent loose stools, magnesium depletion may be severe. This is also true in celiac disease (*gluten enteropathy*) and in those who have had bowel resection surgery. Some evidence suggests vitamin D deficiency may exacerbate magnesium deficiency.¹⁶

Unfortunately, blood magnesium levels are a poor barometer of true body (intracellular) magnesium levels. Only 1% of the body's magnesium is in the blood; the remaining 99% is stored in various body tissues, particularly bone and muscle.¹⁶ If blood magnesium is low, cellular magnesium levels are indeed low—*very* low. If blood magnesium is *normal*, cellular or tissue levels of magnesium may *still* be low. Unfortunately, tissue magnesium levels are not easy to ascertain in living, breathing humans. In one study, only 8% of coronary patients had low blood magnesium, while tissue levels were reduced in 53%.¹⁰ In other words, normal blood magnesium levels do not rule out the possibility of a magnesium deficiency.




Is Your Bottled Water Killing You? *What You Need to Know*

- The mineral magnesium is critical to human health, helping the body perform functions as varied as blood pressure regulation, muscle contraction, heart rhythm stabilization, and nervous system communication.
- Modern water-purification techniques, consumption of magnesium-depleted bottled waters, and poor dietary habits contribute to widespread magnesium deficiencies in the US and around the world. Although the US RDA for magnesium is 420 mg per day for adult men and 320 mg for adult women, most Americans ingest about 270 mg of magnesium per day.
- Magnesium deficiencies have been linked to sudden death from cardiac arrhythmias, along with conditions such as muscle cramps, migraine headaches, and metabolic syndrome.
- Optimal magnesium intake, by contrast, is associated with improved insulin sensitivity, optimal lipoprotein levels, suppression of abnormal heart rhythms, reduced blood pressure, relief from migraine headaches, and improved exercise capability, among other health benefits.
- Strategies for optimizing daily magnesium intake include consuming foods high in magnesium content, supplementing with the soluble fiber known as *inulin*, drinking magnesium-rich mineral waters, and taking magnesium supplements that supply standardized amounts of elemental magnesium.



Short of performing a biopsy to measure tissue magnesium levels, potential magnesium deficiency may manifest as:

- **Cardiac arrhythmias.** Low magnesium levels can trigger abnormal heart rhythms. If you have been diagnosed with a cardiac arrhythmia, it is a good idea to ask your doctor about your magnesium level.¹⁶
- **Low potassium.** Low magnesium commonly accompanies low potassium.¹⁰ Potassium is an electrolyte that is depleted by the use of diuretics and is commonly deficient in many conditions (such as excessive alcohol use, hypertension, and diarrhea). Like magnesium, deficient potassium may not be easily remedied through diet.
- **Muscle cramps.** Magnesium regulates muscle contraction. Leg cramps, or a "charley horse"—painful, vise-like cramps in calves or other muscles—are a common symptom of magnesium deficiency. (Note that leg cramps that occur with physical activity, such as walking, are usually due to atherosclerotic blockages in the leg or abdominal arteries, not low magnesium.)^{17,18}
- **Migraine headaches.** Reflecting magnesium's role in regulating blood vessel tone, low magnesium can trigger vascular spasm in the membranes that cover the brain and spinal cord.¹⁹
- **Metabolic syndrome.** The risk of developing metabolic syndrome—the cluster of metabolic abnormalities that includes low high-density lipoprotein (HDL), high triglycerides, high blood pressure, increased blood sugar, and excessive abdominal fat—is clearly linked with lower magnesium levels. Magnesium plays a fundamental role in regulating cellular responses to insulin.²⁰



The absence of any of these telltale signs does *not* necessarily mean that tissue levels of magnesium are normal. Then how can one know whether magnesium levels are deficient? There is no easy, available method to gauge body magnesium. In all practicality, because of magnesium's crucial role in health, its widespread deficiency in Americans, and the growing depletion of magnesium in water, supplementing with magnesium may be the best way to ensure adequate total body magnesium levels.

Health Benefits of Magnesium Replacement

What can you expect from supplementing magnesium to optimal levels? Research over the past 20 years suggests that magnesium supplementation will accomplish several critically important goals:

- **Magnesium improves insulin sensitivity.** Magnesium helps correct impaired insulin sensitivity, the fundamental defect that characterizes pre-diabetes and metabolic syndrome. An intracellular enzyme called *tyrosine kinase* requires magnesium to allow insulin to exert its blood-sugar-lowering

effects. In several studies, daily oral magnesium supplementation substantially improved insulin sensitivity by **10%** and reduced blood sugar by **37%**.²¹⁻²³

- **Magnesium helps correct abnormal lipoprotein patterns.** Improved insulin sensitivity from magnesium replacement can markedly reduce triglyceride levels.²³ Reduced triglyceride availability, in turn, reduces triglyceride-rich particles, such as very low-density lipoprotein (VLDL) and small low-density lipoprotein (small LDL), both of which are powerful contributors to heart disease.²⁴ Magnesium supplementation can also raise levels of beneficial high-density lipoprotein (HDL).²⁴
- **Magnesium suppresses abnormal heart rhythms.** Magnesium has gained a foothold in hospital care following coronary bypass surgery, when the abnormal heart rhythm known as *atrial fibrillation* commonly occurs. Magnesium may help suppress this rhythm outside of the hospital as well,²⁵ suggesting a preventive role in averting abnormal heart rhythms.

- **Magnesium reduces blood pressure.** Magnesium regulates blood pressure by modulating vascular tone. Magnesium works in ways similar to the prescription hypertension drugs known as *calcium-channel antagonists* (such as *diltiazem* and *nifedipine*), which block calcium channels that trigger constriction of the arteries. Magnesium stimulates the production of *prostacyclins* and *nitric oxide*, which are potent artery-relaxing agents.²⁶ Magnesium exerts a modest effect of reducing blood pressure, reflecting its whole-body artery-relaxing properties.²⁷
- **Magnesium can block migraine headaches.** Magnesium has been explored as a means to prevent or relieve migraine headaches. People suffering migraine headaches tend to have lower magnesium levels.²⁸ A study from the State University of New York showed that intravenous magnesium relieved headache symptoms in 15 minutes in 80% of recipients.²⁹ Other studies have since corroborated magnesium's beneficial effect on migraine headaches, including a trial in children, in which *oral* supplementation with magnesium oxide reduced the frequency and severity of migraine.³⁰
- **Magnesium may improve exercise performance.** Extensive research in athletes has found that intensive exercise triggers magnesium loss through urinary excretion and perspiration. When magnesium is low, supplementation enhances exercise performance by reducing lactate blood levels (indicating brief, strength-based anaerobic muscle activity), decreasing oxygen requirements, and increasing muscle strength.^{31,32}
- **Magnesium may benefit many other conditions.** Other conditions in which magnesium

is believed to exert positive effects include fibromyalgia,³³ asthma (acute episodes have been treated successfully with both intravenous and aerosolized magnesium),³⁴ prevention of osteoporosis,³⁵ and premenstrual syndrome.³⁶

Can you correct metabolic syndrome and its complications—such as insulin resistance and high blood pressure—without replacing magnesium? Of course you can, just as you can operate your car without changing the oil. However, magnesium deficiency will catch up with you, and consuming this basic supplement will help you to more easily achieve your health goals.

Strategies for Optimizing Your Magnesium Intake

According to the US Department of Health and Human Services, nearly all of us fail to achieve even the modest magnesium RDAs of 420 mg for adult males and 320 mg for adult females. Most American adults ingest about 270 mg of magnesium a day, well below the RDA and enough to generate a substantial cumulative deficiency over months and years.³⁷

The magnesium RDA refers to *elemental* magnesium, defined as the amount of magnesium regardless of its source or form. Magnesium is gen-

erally available as various "salts" (not to be confused with table salt), and the amount of elemental magnesium contained in each varies depending on the salt. For example, the amount of magnesium in magnesium oxide is 60%; in magnesium carbonate, 45%; in magnesium citrate, 16%; and in magnesium chloride, 12%.³⁸ Thus, magnesium oxide supplements tend to contain more elemental magnesium per pill than do magnesium citrate supplements.

Magnesium salts differ in absorption. Magnesium oxide, though inexpensive and widely available, is thought to be relatively less absorbed than the citrate and chloride forms.³⁹⁻⁴¹

You can also increase your magnesium intake by choosing foods rich in magnesium, which are listed in the table below.⁴²

Dietary Sources of Magnesium

Nuts, pumpkin seeds, spinach, and oat bran are particularly rich and healthy sources of magnesium.

Another strategy for boosting magnesium intake is to supplement your diet with the soluble fiber known as *inulin*. Like other soluble fibers, inulin may exert modest cholesterol- and triglyceride-reducing effects. However, it also enhances magnesium absorption in the intestine.⁴³ Inulin can be taken as a supplement,

FOODS RICH IN MAGNESIUM (magnesium content in mg)

Almonds (1 oz; 24 nuts)	78	Oatmeal (1 cup, cooked)	56
Artichokes (1 cup)	101	Pumpkin seeds (1 oz; 142 seeds)	151
Barley (1 cup, raw)	158	Rice, brown (1 cup, cooked)	84
Beans, black (1 cup, cooked)	120	Soybeans (1 cup, cooked)	148
Beans, lima (1 cup, cooked)	101	Spinach (1 cup, cooked)	163
Brazil nuts (1 oz; 6-8 nuts)	107	Trail mix (1 cup)	235
Halibut (1/2 filet)	170	Walnuts (1 oz; 14 halves)	45
Filberts, hazelnuts (1 oz)	46	Wheat flour, whole grain (1 cup)	166
Oat bran (1 cup, raw)	221		

Source: USDA National Nutrient database for Standard Reference, Release 31

MAGNESIUM CONTENT OF MINERAL WATERS

The following waters contain far more than the usual amounts of magnesium. Some, like Apollinaris and Pellegrino, are widely available in American grocery stores, while others are found only in upscale groceries or through websites of the water producers.

Mineral Water	Magnesium Content
Original Fountain of Youth Mineral Water (Florida)	609 mg/L
Apollinaris (Germany) (410 mg/L of sodium)	130 mg/L
Adobe Springs (California and other western states)	110 mg/L
Badoit (France)	85 mg/L
Colfax (Iowa)	91 mg/L
Deep Rock (Colorado)	60 mg/L
Evian	24 mg/L
Gerolsteiner (Germany)	108 mg/L
Noah's California Spring Water	110 mg/L
Pellegrino Sparkling Mineral Water (Italy) (43.6 mg/L of sodium)	55.9 mg/L
Manitou Mineral Water (Colorado)	43 mg/L
Rosbacher	93 mg/L
St. Gero	109.4 mg/L

Both Apollinaris and Pellegrino contain more sodium than most other waters, and therefore should be avoided by those who are limiting their sodium intake due to existing hypertension, fluid retention, or kidney disease.

and is contained in some foods (for example, the Stonyfield Farms brand of yogurt). Inulin can help increase satiety (the sense of fullness you get with eating), resulting in decreased calorie intake throughout the day.⁴⁴ Inulin thus holds promise in supporting efforts to lose weight.⁴⁵

One more important way to optimize your magnesium intake is to choose water that is rich in magnesium. Unfortunately, in the US, this is easier said than done. The FDA regulates bottled water and mandates that the only additives permitted are fluoride and antimicrobials to deter bacterial growth. Magnesium cannot therefore be added to water labeled spring water or mineral water.

Magnesium-rich mineral waters are not easy to find, but they are out there. By FDA definition, mineral waters must contain at least 250 parts per million (ppm) of total dissolved solids. Not all mineral water contains significant quantities of magnesium. For example, Napa Valley's Calistoga Springs, labeled as "mineral water," contains 0.61–0.96 mg/L of magnesium, or virtually none.

To determine the amount of magnesium contained in bottled water labeled "mineral water" but not listed above, go to the bottler's website to determine the water's composition.

With the exception of Florida's Original Fountain of Youth Mineral Water, drinking an entire liter of many so-called mineral waters provides only a modest amount of magnesium. Thus, for instance, if you are currently ingesting around 250 mg a day of magnesium from your diet, drinking a liter

of Gerolsteiner a day (supplying 108 mg/L of magnesium) will increase your magnesium consumption only to about 350 mg per day. However, by adding a magnesium supplement that provides as little as 100 mg of elemental magnesium, you will have more than achieved the RDA for an adult male. Since many mineral waters are expensive (around \$2-3 per liter), magnesium supplements are a far less costly way to optimize your magnesium intake.

Conclusion

The intensification of municipal water treatment has resulted in a growing epidemic of magnesium deficiency, with most Americans falling even to achieve the modest levels set by the government-recommended RDA. Most of us have daily deficiencies in magnesium intake of only 70–200 mg a day.

The consequences of magnesium deficiency can be dramatic, including poor insulin response, migraine headaches, high blood pressure, and abnormal and even dangerous heart rhythms.

Fortunately, there are plenty of healthy choices—foods rich in magnesium, low-cost magnesium supplements, and waters rich in magnesium—that can you help reach or exceed the magnesium RDA and attain the numerous health benefits conferred by optimal magnesium intake. ■

MAGNESIUM DOSAGE GUIDELINES

- The recommended dietary allowance (RDA) for magnesium is 420 mg a day for adult men and 320 mg a day for adult women.⁴⁶ Most people fail to achieve the RDA, which may lead to magnesium deficiency.⁴⁷
- The most common adverse reaction from the use of magnesium supplements is diarrhea. Other gastrointestinal symptoms include nausea and abdominal cramping. Diarrhea and other gastrointestinal symptoms are less likely to occur if magnesium supplements are taken with food.⁴⁸
- Magnesium supplements are contraindicated in those with kidney failure. Those with *myasthenia gravis* (an autoimmune disorder that results in progressive skeletal muscle weakness) should avoid magnesium supplements.⁴⁶

Dr. William Davis is an author and cardiologist practicing in Milwaukee, WI. He is founder of the *Track Your Plaque* program, a heart disease prevention and reversal program that shows how CT heart scans can be used to track and control coronary plaque. He can be reached at www.TrackYourPlaque.com.

References

1. Available at: <http://www.historyofwaterfilters.com/>. Accessed November 14, 2006.
2. Azoulay A, Garzon P, Eisenberg MJ. Comparison of the mineral content of tap water and bottled waters. *J Gen Intern Med*. 2001 Mar;16(3):160-75.
3. Available at: <http://api.oregonstate.edu/infocenter/minerals/magnesium/index.html>. Accessed November 14, 2006.
4. Touyz RM. Magnesium in clinical medicine. *Front Biosci*. 2004 May 19;9:278-93.
5. Liebscher DH, Liebscher DE. About the misdiagnosis of magnesium deficiency. *J Am Coll Nutr*. 2004 Dec;23(6):730S-1S.
6. Durlach J, Bac P, Durlach V, et al. Magnesium status and ageing: an update. *Magnes Res*. 1998 Mar;11(1):25-42.
7. Eisenberg MJ. Magnesium deficiency and sudden death. *Am Heart J*. 1992 Aug;124(2):544-9.
8. Dacey MJ. Hypomagnesemic disorders. *Crit Care Clin*. 2001 Jan;17(1):155-73, viii.
9. Eisenberg MJ. Magnesium deficiency and cardiac arrhythmias. *NY State J Med*. 1986 Mar;86(3):133-6.
10. Purvis JR, Movahed A. Magnesium disorders and cardiovascular diseases. *Clin Cardiol*. 1992 Aug;15(8):556-68.
11. Smetana R, Stuhlinger HG, Kiss K, Glogar DH. Intravenous magnesium sulphate in acute myocardial infarction—is the answer “MAGIC”? *Magnes Res*. 2003 Mar;16(1):65-9.
12. Kousa A, Havulinna AS, Molchanova E, et al. Calcium-magnesium ratio in local groundwater and incidence of acute myocardial infarction among males in rural Finland. *Environ Health Perspect*. 2006 May;114(5):730-4.
13. Anderson TW, Le Riche WH, MacKay JS. Sudden death and ischemic heart disease. Correlation with hardness of local water supply. *N Engl J Med*. 1969 Apr 10;280(15):805-7.
14. Available at: http://www.nsf.org/international/press_release.asp?p_id=12041. Accessed November 14, 2006.
15. Laires MJ, Monteiro CP, Bicho M. Role of cellular magnesium in health and human disease. *Front Biosci*. 2004 Jan 19;9:262-76.
16. Berkelhammer C, Bear RA. A clinical approach to common electrolyte problems: 4. Hypomagnesemia. *Can Med Assoc J*. 1985 Feb 15;132(4):360-8.
17. Roffe C, Sills S, Crome P, Jones P. Randomised, cross-over, placebo controlled trial of magnesium citrate in the treatment of chronic persistent leg cramps. *Med Sci Monit*. 2002 May;8(5):CR326-30.
18. Bilbey DL, Prabhakaran VM. Muscle cramps

THE BASICS OF WATER AND WATER PURIFICATION

While the Environmental Protection Agency regulates the quality of tap water, the Food and Drug Administration (FDA) is responsible for regulating bottled water. In 1995, the FDA issued its most recent regulations classifying various waters:

- **Artesian well water** is water that naturally flows upward from an underground aquifer to a well, without the need for pumping.
- **Mineral water** is water from an underground source that contains at least 250 parts per million (ppm) of dissolved solids consisting of minerals and trace elements. Mineral content of 250-500 ppm is often called “low mineral content” or “light mineral water,” while content of 1500 ppm or greater is “high mineral content.” (In Europe, spring waters with dissolved solids equal to or less than 500 mg/L are considered “mineral with low mineral content” or simply “mineral water.”) Minerals and trace elements cannot be added artificially to water labeled as mineral.
- **Spring water**, like artesian well water, comes from an underground source but flows naturally to the earth’s surface. It cannot come from a public or municipal source. Spring water must be collected directly at the spring or through a borehole tapping the underground source. Mineral content is less than 250 ppm and cannot be added after collection.
- **Well water** is water from a hole bored or drilled into the ground, which taps into an aquifer and is drawn to the surface using a pump. Many homes in the US that do not have access to municipal water use well water.

To make matters even more complicated, any water—regardless of the source—can be treated or filtered. This is usually done to modify its taste or to remove undesirable ingredients. Methods of treatment are defined as:

- **Distillation.** Water is vaporized and collected, leaving behind any solid residues, including minerals. Distilled water contains no minerals whatsoever.
- **Reverse osmosis.** In this common water-purifying process, water is forced through membranes to remove minerals in the water.
- **Deionization.** Also called *demineralization* or *ion exchange*, this process uses synthetic resins to remove ions and minerals from water. This is very effective at removing ionized impurities, but does not remove organic, bacterial, pathogenic, or particulate matter efficiently. Deionized water contains no magnesium.
- **Absolute 1 micron filtration.** Water is passed through filters that remove particles larger than 1 micron in size, including *Cryptosporidium*, a parasite that causes intestinal infestation. This process does not affect the water’s mineral content.
- **Ozonation.** Many bottled water companies use this process instead of chlorine to rid water of bacteria. Ozonation does not affect the mineral content of water.

Many bottled waters are simply tap water processed using one or more of the above processes of distillation, reverse osmosis, deionization, or filtration. This leaves the water virtually devoid of both nutrients and contaminants. Of the 700 or so brands of bottled water available in the US, 80% are processed water. Many experts say that treated water like this is virtually identical to that produced by home water purifiers. The appeal of these waters is therefore a reduction in impurities like lead and pesticide residues, or better taste—but not enhanced mineral content. Bottled processed waters contain *little or no* magnesium.

It should also be noted that unlike tap water, purified waters and water purifiers reduce or eliminate the fluoride that is added by many municipal treatment facilities to promote dental health. Although the FDA permits producers to add it back to purified water, few actually do.

Waters derived from natural sources like artesian well water, well water, mineral water, and spring water are generally *slightly* richer in mineral content than are processed and tap waters. However, the difference is small. Nearly all American bottled waters obtained from natural sources—whether artesian, well, spring, or mineral waters—contain less than 6 ppm of magnesium, a trivial amount.

A ONE-MAN CRUSADE FOR MAGNESIUM SUPPLEMENTATION

A year and a half ago, we called attention to the remarkable efforts of Paul Mason, a Life Extension member who has painstakingly documented that magnesium deficiency is a significant cause of cardiovascular disease ("How Many Americans Are Magnesium Deficient? *Life Extension*, September 2005). Paul has made it his personal mission to persuade the world that consuming more magnesium would prevent tens of millions of premature deaths from heart attack, stroke, and other diseases.

Paul took the extraordinary step of suing the FDA and Justice Department to mandate that all bottled water and soft drinks be fortified with magnesium. He contends that adding magnesium to bottled drinks would provide most Americans with the government's recommended daily intake of this lifesaving mineral. Paul sought to have a federal court order the FDA to inform the public about the critical need to ingest adequate magnesium.

To substantiate his position, Paul Mason created a website (www.mgwater.com) providing links to hundreds of published scientific papers documenting how magnesium prevents heart attacks and other life-threatening health conditions. He has collected 80 studies showing that small amounts of magnesium in water, often just 5-20 mg per liter, lower incidences of mortality and heart disease.

According to data compiled by Paul Mason, the US National Academy of Sciences has reported that more than 50 studies from nine countries "have indicated an inverse relationship between water hardness and mortality from cardiovascular disease. That is, people who drink water that is deficient in magnesium and calcium generally appear more susceptible to this disease. The US National Academy of Sciences has estimated that a nationwide initiative to add calcium and magnesium to soft water might reduce the annual cardiovascular death rate by 150,000 in the United States."

Despite being rebuffed by the US government, Paul Mason continues to work with the World Health Organization to encourage adding magnesium to drinking water in other countries whose citizens suffer from diseases related to magnesium deficiency. As he reports on his website, the potential health benefits of such initiatives are nothing less than staggering:

"Even modest amounts of magnesium in water apparently reduce the rate of mortality (by an average of .003345). Applying that mathematical benchmark average to a world population of 6,350,000,000 suggests a possible decline in mortality of 21,240,750 per year—making magnesium deficiency a bigger cause of death than AIDS and war combined."

To learn more, please visit Paul Mason's website at www.mgwater.com

- and magnesium deficiency: case reports. *Can Fam Physician*. 1996 Jul;42:1348-51.
- Bussone G. Pathophysiology of migraine. *Neurol Sci*. 2004 Oct;25 Suppl 3S239-41.
 - He K, Liu K, Davighis ML, et al. Magnesium intake and incidence of metabolic syndrome among young adults. *Circulation*. 2006 Apr 4;113(13):1675-82.
 - Guerrero-Romero E, Tamez-Perez HE, Gonzalez-Gonzalez G et al. Oral magnesium supplementation improves insulin sensitivity in non-diabetic subjects with insulin resistance. A double-blind placebo-controlled randomized trial. *Diabetes Metab*. 2004 Jun;30(3):253-8.
 - Rodriguez-Moran M and Guerrero-Romero E. Oral magnesium supplementation improves insulin sensitivity and metabolic control in type 2 diabetic subjects: a randomized double-blind controlled trial. *Diabetes Care*. 2003 Apr;26(4):1147-52.
 - Yokota K, Kato M, Lister E et al. Clinical efficacy of magnesium supplementation in patients with type 2 diabetes. *J Am Coll Nutr*. 2004 Oct;23(5):506S-9S.
 - Rasmussen HS, Aurup P, Goldstein K, et al. Influence of magnesium substitution therapy on blood lipid composition in patients with ischemic heart disease. A double-blind, placebo controlled study. *Arch Intern Med*. 1989 May;149(5):1050-3.
 - Piotrowski AA, Kalus JS. Magnesium for the treatment and prevention of atrial tachyarrhythmias. *Pharmacotherapy*. 2004 Jul;24(7):879-95.
 - Sontia B, Touyz RM. Role of magnesium in hypertension. *Arch Biochem Biophys*. 2006 May 24.
 - Jee SH, Miller ER, III, Guallar E, et al. The effect of magnesium supplementation on blood pressure: a meta-analysis of randomized clinical trials. *Am J Hypertens*. 2002 Aug;15(8):691-6.
 - Gallai V, Sarchielli F, Morucci P, Abbritti G. Magnesium content of mononuclear blood cells in migraine patients. *Headache*. 1994 Mar;34(3):160-5.
 - Mauskop A, Altrura BT, Cracco RQ, Altura BM. Intravenous magnesium sulfate rapidly alleviates headaches of various types. *Headache*. 1996 Mar;36(3):154-60.
 - Wang F, Van Den Eeden SK, Ackerson LM, et al. Oral magnesium oxide prophylaxis of frequent migrainous headache in children: a randomized, double-blind, placebo-controlled trial. *Headache*. 2003 Jun;43(6):801-10.
 - Lukaski HC. Magnesium, zinc, and chromium nutrition and athletic performance. *Can J Appl Physiol*. 2001;26 SupplS13-S22.
 - Cinar V, Nizamlioglu M, Mogulkoc R. The effect of magnesium supplementation on lactate levels of sportsmen and sedenter. *Acta Physiol Hung*. 2006 Jun;93(2-3):137-44.
 - Sarac AJ, Gur A. Complementary and alternative medical therapies in fibromyalgia. *Curr Pharm Des*. 2006;12(1):47-57.
 - Blitz M, Blitz S, Hughes R, et al. Aerosolized magnesium sulfate for acute asthma: a systematic review. *Chest*. 2005 Jul;128(1):337-44.
 - Rude RK, Gruber HE. Magnesium deficiency and osteoporosis: animal and human observations. *J Nutr Biochem*. 2004 Dec;15(12):710-6.
 - Rappin A. A review of treatment of premenstrual syndrome and premenstrual dysphoric disorder. *Psychoneuroendocrinology*. 2003 Aug;28 Suppl 3:39-53.
 - Ervin RB, Wang CY, Wright JD, Kennedy-Stephenson J. Dietary intake of selected minerals for the United States population: 1989-2000. *Adv Data*. 2004 Apr 27;(341):1-5.
 - Available at: <http://ods.od.nih.gov/factsheets/magnesium.asp#6>. Accessed November 15, 2006.
 - Firooz M, Graber M. Bioavailability of US commercial magnesium preparations. *Magnes Res*. 2001 Dec;14(4):257-62.
 - Coudray C, Rambreau M, Feillet-Coudray C, et al. Study of magnesium bioavailability from ten organic and inorganic Mg salts in Mg-depleted rats using a stable isotope approach. *Magnes Res*. 2005 Dec;18(4):215-23.
 - Walker AE, Marakis G, Christie S, Byng M. Mg citrate found more bioavailable than other Mg preparations in a randomized, double-blind study. *Magnes Res*. 2003 Sep;16(3):183-91.
 - Available at: <http://www.nal.usda.gov/fnic/foodcomp/Data/SR14/wtrank/sr14w304.pdf>. Accessed November 15, 2006.
 - Roberfroid MB. Introducing inulin-type fructans. *Br J Nutr*. 2005 Apr;93 Suppl 1S13-S25.
 - Archer BJ, Johnson SK, Devereux HM, Baxter AL. Effect of fat replacement by inulin or lupin-kernel fibre on sausage patty acceptability, post-meal perceptions of satiety and food intake in men. *Br J Nutr*. 2004 Apr;91(4):591-9.
 - Hoeger WW, Harris C, Long EM, Hopkins DR. Four-week supplementation with a natural dietary compound produces favorable changes in body composition. *Adv Ther*. 1998 Sep-Oct;15(5):305-14.
 - Available at: http://www.pdrhealth.com/drug_info/nmdrugprofiles/nutsupdrugs/mag_0167.shtml. Accessed November 16, 2006.